

University of Groningen

The therapeutic potential of adenoviral gene therapy and angiotensine-(1-7) in proteinuric kidney disease

Wouden, Esther Anita van der

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:
2007

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Wouden, E. A. V. D. (2007). *The therapeutic potential of adenoviral gene therapy and angiotensine-(1-7) in proteinuric kidney disease*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

**The therapeutic potential of
adenoviral gene therapy and angiotensin-(1-7)
in proteinuric kidney disease**

Els van der Wouden

The publication of this thesis was financially supported by:
Dutch Kidney Foundation
Rijks*universiteit* Groningen
Groningen University Institute for Drug Exploration (GUIDE)

CIP-gegevens koninklijke bibliotheek, Den Haag

Van der Wouden, E.A.

The therapeutic potential of adenoviral gene therapy and angiotensin-(1-7) in proteinuric kidney disease

Proefschrift Groningen – met literatuur opgave – met samenvatting in het Nederlands

ISBN 9789036730419

© Copyright 2007 E.A. van der Wouden

All rights are reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, mechanically, by photocopying, recording, or otherwise, without the written permission of the author.

© Cover design: Daniël van der Veer

Printed by: Gildeprint Drukkerijen BV, Enschede

Rijksuniversiteit Groningen

**The therapeutic potential of
adenoviral gene therapy and angiotensin-(1-7)
in proteinuric kidney disease**

Proefschrift

ter verkrijging van het doctoraat in de
Medische Wetenschappen
aan de Rijksuniversiteit Groningen
op gezag van de
Rector Magnificus, dr. F. Zwarts,
in het openbaar te verdedigen op
woensdag 23 mei 2007
om 13:15 uur

door

Esther Anita van der Wouden
geboren op 25 november 1977
te Staphorst

Promotores: Prof. dr. R.H. Henning
Prof. dr. D. de Zeeuw

Copromotor: Dr. L.E. Deelman

Beoordelingscommissie: Prof. dr. H.J. Haisma
Prof. dr. G.J. Navis
Prof. dr. K. Sharma

Paranimfen:

P.E. Ettema – van der Wouden

Dr. M. Sandovici

Contents

1. Introduction and aim of the thesis	9
2. Approaches and methods in gene therapy for kidney disease <i>J Pharmacol Toxicol Methods.</i> 2004; 50:13-24	15
3. Adenovirus harbouring an RGD-motif in the HI-loop is a fibroblast-specific vector in healthy and nephrotic kidneys	37
4. The role of angiotensin-(1-7) in renal vasculature of the rat <i>J Hypertens.</i> 2006; 24:1971-8	53
5. Recombinant adenovirus reduces proteinuria in adriamycin-induced nephrosis in the rat	67
6. Does angiotensin-(1-7) contribute to the antiproteinuric effect of ACE inhibitors? <i>J Renin Angiotensin Aldosterone Syst.</i> 2005; 6:96-101	81
7. Summary, discussion and future directions	93
Samenvatting	101
<i>Appendices</i>	
8. Altered dosing of trandolapril does not overcome nocturnal resistance to the antiproteinuric effect of ACE inhibition in nondiabetic kidney disease	109
9. TGF- β 1 inhibits angiotensin II-induced MAP kinase signalling in aortic smooth muscle cells through downregulation of angiotensin II type 1 receptors	121
Dankwoord	131

